

Equivalent solutions may exist

Problem 1

17 marks

Thermal and photolytic decomposition of Acetaldehyde

1.1 a)

$$-\frac{d[\text{CH}_3\text{CHO}]}{dt} = k[\text{CH}_3\text{CHO}]^{3/2}$$


$$\frac{d[\text{CH}_4]}{dt} = \frac{d[\text{CO}]}{dt} = k[\text{CH}_3\text{CHO}]^{3/2}$$

(b)

$$\text{Order} = 3/2$$

$$\text{Rate} = 8 v$$

1.2



1.3 (a)

Propogation steps : (ii) and (iii)
Termination step: (iv)

(b)


$$[\bullet\text{CH}_3] = \left(\frac{k_1}{2k_4}\right)^{1/2} [\text{CH}_3\text{CHO}]^{1/2}$$

$$[\bullet\text{CH}_3\text{CO}] = \frac{k_2}{k_3} \left(\frac{k_1}{2k_4}\right)^{1/2} [\text{CH}_3\text{CHO}]^{3/2}$$

(c)

$$\frac{d[\text{CO}]}{dt} = k_2 \left(\frac{k_1}{2k_4}\right)^{1/2} [\text{CH}_3\text{CHO}]^{3/2}$$

1.4 (a)



$$\text{Rate} = I_{\text{abs}}$$

$$[\bullet\text{CH}_3] = (I_{\text{abs}}/2k_4)^{1/2} [\text{CH}_3\text{CHO}]^{3/2}$$

$$d[\text{CO}]/dt = k_2 \times (I_{\text{abs}}/2k_4)^{1/2} [\text{CH}_3\text{CHO}]^{3/2}$$

(b)

$$\lambda = 317.5 \text{ nm}$$

1.5 (a)

$$E_{\text{thermal}} = E_2 + \frac{1}{2}(E_1 - E_4)$$

(b)

$$E_{\text{photochemical}} = E_2 - \frac{1}{2} E_4$$

1.6


(a)

$$E_{\text{thermal}} = 196.46 \text{ kJ mol}^{-1}$$

(b)

$$5.879 \times 10^{-4} \text{ mol dm}^{-3} \text{ sec}^{-1}$$

1.7 (a)



(b)


$$[\text{CH}_3\text{CHO}] = k_5/k_6 [\text{CH}_3\text{CH}_2\text{OH}]$$

Problem 2



19 marks

Chemistry of coordination compounds

2.1



or balanced equation with $[\text{Cu}(\text{H}_2\text{O})_4]^{2+}$ entity



2.2

b] completely filled d-level in Cu(I)


2.3

a] oxidation state of the metal.

b] nature of the ligand.

c] geometry of the complex.

2.4



$[\text{Ni}(\text{H}_2\text{O})_6]^{2+}$ is octahedral.

$[\text{NiCl}_4]^{2-}$ is tetrahedral

2.5

- a]. IUPAC Name : Dichlorobis(ethylenediamine)cobalt(III) ion.
 Dichlorobis(ethane-1,2-diamine)cobalt(III) ion
 Dichloridobis(ethylenediamine)cobalt(III) ion.
 Dichloridobis(ethane-1,2-diamine)cobalt(III) ion

b]. Geometrical isomers:



c] cis-[CoCl₂(en)₂]⁺ is optically active.

d] Two optical isomers of cis-[CoCl₂(en)₂]⁺ :



2.6

Ni²⁺ [Ar] 3d⁸



[NiCl₄]²⁻



sp³

sp³ hybridization

Tetrahedral

Paramagnetic (2 unpaired electrons)

Pt²⁺ [Xe] 5d⁸



[PtCl₄]²⁻



dsp²

dsp² hybridization


Square planar

Diamagnetic (no unpaired electrons)

2.7

A	B Δ_o (cm ⁻¹)
i) $[\text{CrF}_6]^{3-}$	d) 15,000
ii) $[\text{Cr}(\text{H}_2\text{O})_6]^{3+}$	c) 17,400
iii) $[\text{CrF}_6]^{2-}$	b) 22,000
iv) $[\text{Cr}(\text{CN})_6]^{3-}$	a) 26,600

2.8

Answer:


Oxidation state

Fe(III)

Ni(0)

Coordination No.

of Fe(III) : 6


of Ni(II) : 4

EAN of central metal ion

35

36





2.9



a) (i) by elongation along z-axis.

 b) (ii) dx^2-y^2 orbital.

2.10




Problem 3


14 marks

Chemistry of isomeric benzenes


3.1




3.2




3.3



3.4



3.5





3.6



(b) Three

X


3.7




3.8



3.9



3.10




Problem 4

10 marks



s-Block Elements

4.1 a) only one valence electron

b) large atomic size



4.3




4.4


i) Bond order = 1

ii) diamagnetic

4.5



OR




- 4.6 b) It is paramagnetic in nature X
 c) On standing this solution slowly liberates hydrogen
 resulting in the formation of sodium amide X
- 4.7 c) half the number of tetrahedral X
- 4.8 b) cyclohexane X
 c) diisopropyl ether X
- 4.9 a) ionization energy of alkali metal X
 b) electron gain enthalpy of halogen X
 d) sizes of cations and anions X
- 4.10

Problem 5

17 marks

Carboxylic acid derivatives

5.1



5.2 (c) Amide > Ester > Acid Chloride

X

5.3 Amide

X

5.4

1650 cm^{-1}	A
1750 cm^{-1}	C
1800 cm^{-1}	B

5.5 $\text{CH}_3\text{CH}_2\text{COCl}$

X



5.6 Best

C



Poorest

B




5.7




5.8



5.9



5.10




5.11 (i)

9


(ii)

Both are S



5.12



5.13



5.14



Problem 6

17 marks

Chemical Thermodynamics

6.1

$$2930 \text{ J}$$

6.2

$$K_p = 0.7030$$

$$K_p = K_c$$

6.3

$$X_{\text{CO}} = 0.342, X_{\text{H}_2} = 0.458, X_{\text{H}_2\text{O}} = 0.092, X_{\text{CO}_2} = 0.108$$

$$X_{\text{CO}} = 34.95\%, X_{\text{H}_2} = 45.41\%, X_{\text{H}_2\text{O}} = 9.59\%, X_{\text{CO}_2} = 10.06\%$$

6.4

$$\Delta H_{1400} = 31258 \text{ J}$$

6.5

a) K_p will increase with increase in temperature

6.6

$$\text{Air intake (engine; m}^3\text{s}^{-1}) = V_A = 4 \times 9.902 \times 10^{-3} \text{ m}^3\text{s}^{-1} = 0.0396 \text{ m}^3\text{s}^{-1}$$

6.7

$$T_1 = 2060 \text{ K}$$

$$T_2 = 708 \text{ K}$$

6.8

Compound	Molar composition of gases after leaving the bed (Mol $\times 10^{-4}$)
N_2 (g)	407.64
O_2 (g)	38.55
CO (g)	0.78
CO_2 (g)	44.14
H_2O (g)	49.44


Problem 7

10 marks

7.1

$$V = 87.5 \text{ mL}$$

7.2



$$K_a = \frac{[\text{H}_3\text{O}^+][\text{CH}_3\text{COO}^-]}{[\text{CH}_3\text{COOH}]}$$

7.3


$$[\text{H}_3\text{O}^+] = -\frac{K_a \pm \sqrt{K_a^2 + 4K_a C_T}}{2}$$

7.4

$$\text{pH} = 2.88$$

7.5

a)



b)

$$[\text{CH}_3\text{COOH}]_{\text{eq}} = C_T - [\text{H}_3\text{O}^+] + [\text{OH}^-]$$

$$[\text{CH}_3\text{COO}^-]_{\text{eq}} = [\text{CH}_3\text{COONa}] + [\text{H}_3\text{O}^+] - [\text{OH}^-]$$

c)

$$[\text{H}_3\text{O}^+] = K_a \frac{C_T}{[\text{CH}_3\text{COONa}]}$$

d)

$$\text{pH} = 3.80$$

7.6

$$\text{pH} = 8.73$$