

P1₁

Name: _____

P1₁

Student code: _____

1.1 You started with 2.54 g of polycarbonate. Determine the theoretical yield of bisphenol A in g. (2 points)

$$M_1 (\text{polycarbonate}) = M_1 (\text{C}_{16}\text{H}_{14}\text{O}_3)_n\text{H}_2 \approx M_1 (\text{C}_{16}\text{H}_{14}\text{O}_3) = 254.30 \text{ g/mol}$$

$$m_1 = 2.54 \text{ g}$$

$$M_2 (\text{C}_{15}\text{H}_{16}\text{O}_2) = 228.31 \text{ g/mol}$$

$$m_2 = m_1 \cdot M_1^{-1} \cdot M_2$$

2.28 g

Theoretical yield of bisphenol A:

exact answer: 2 points; incorrect mathematical rounding, more or less than two figures after the decimal point (e.g. 2.3 g, 2.281 g): 1 point; wrong or missing answer: 0 points.

1.2 Determine your theoretical yield of bisphenol A bis(carboxymethyl)ether in g based on 2.00 g bisphenol A. (2 points)

$$M_2 (\text{C}_{15}\text{H}_{16}\text{O}_2) = 228.31 \text{ g/mol}$$

$$m_2 = 2.00 \text{ g}$$

$$M_3 (\text{C}_{19}\text{H}_{20}\text{O}_6) = 344.39 \text{ g/mol}$$

$$m_3 = m_2 \cdot M_2^{-1} \cdot M_3$$


3.02 g

Theoretical yield of bisphenol A bis(carboxymethyl)ether:

exact answer: 2 points; incorrect mathematical rounding, more or less than two figures after the decimal point (e.g. 3.0 g, 3.017 g): 1 point; wrong or missing answer: 0 points.


1.3 Unwanted by-products are possible in the second step. Write down the structural formulas of two most probable unwanted by-products. (6 points)

1. Bisphenol A reacts only once with sodium chloroacetate (monosubstitution):



(3)

2. Alkaline hydrolysis of sodium chloroacetate:



(3)

For each of the two answers - exact structural formula: 3 points, one careless mistake: 1 point less, two careless mistakes: 2 points less, wrong or missing answers: 0 points.

1.4 Step 1, yield [%] of the product measured by the organizer: (30.0 points)

$$f(x) = 0 \quad x < 61$$


$$f(x) = x - 61 \quad 61 \leq x \leq 91$$

$$f(x) = 30 \quad 91 < x \leq 95 \quad \text{master value}$$

$$f(x) = -3x + 315 \quad 95 < x \leq 105$$

$$f(x) = 0 \quad x > 105$$

$$m_2 \cdot M_1 \cdot m_1^{-1} \cdot M_2^{-1} \cdot 100 = x \quad [\%]$$



1.5 Step 1, melting point [°C] of the product measured by the organizer: (10.0 points)


$$f(x) = 0 \quad x < 146.0$$

$$f(x) = +1.03093x + 150.51546 \quad 146.0 \leq x < 155.7$$

$$f(x) = 10 \quad 155.7 \leq x \leq 156.2 \quad \text{master value}$$

$$f(x) = -5.55556x + 877.77778 \quad 156.2 < x \leq 158.0$$

$$f(x) = 0 \quad x > 158.0$$



1.6 Step 2, yield [%] of the product measured by the organizer: (30.0 points)

$$f(x) = 0 \quad x < 30$$


$$f(x) = 0.75x - 22.5 \quad 30 \leq x \leq 70$$

$$f(x) = 30 \quad 70 < x \leq 80 \quad \text{master value}$$

$$f(x) = -1.5x + 150 \quad 80 < x \leq 100$$

$$f(x) = 0 \quad x > 100$$

$$m_3 \cdot M_2 \cdot m_2^{-1} \cdot M_3^{-1} \cdot 100 = x \quad [\%]$$



1.7 Step 2, melting point [°C] of the product measured by the organizer: (20.0 points)


$$f(x) = 0 \quad x < 162$$

$$f(x) = 2x - 324 \quad 162 \leq x \leq 172$$

$$f(x) = 20 \quad 172 < x \leq 174 \quad \text{master value}$$

$$f(x) = -10x + 1760 \quad 174 < x \leq 176$$

$$f(x) = 0 \quad x > 176$$



A penalty of 10 points will be given if melting point tubes are not filled by the student.

Accuracy of points for 1.4 – 1.7: rounding value is one digit after the decimal point.

P2₁

Name: _____

P2₁

Student code: _____

2.1 Which alkaline earth metal(s) can be found in the superconductor? Mark only one box! (30)

- Ca (0) Sr (0) Ba x (30)
 Ca and Sr (0) Ca and Ba (5) Sr and Ba (15)
 Ca and Sr and Ba (10)

Complete the following reaction equations: (2)**2.2 Quantitative determination of the total content of lanthanum and copper. (35)**



Titration No.	V _{initial} (mL)	V _{final} (mL)	V (mL)
1			
2			
3			
...			
...			
...			

appropriate consumption of 0.1000 mol L⁻¹ EDTA solution V = 11.60*mL
 (according to 100 mL of superconductor solution)

2.3 Quantitative determination of the copper content. (35)

Titration No.	V _{initial} (mL)	V _{final} (mL)	V (mL)
1			
2			
3			
...			
...			
...			

appropriate consumption of 0.01000 mol L⁻¹ Na₂S₂O₃ solution V = 10.50* mL
 (according to 100 mL of superconductor solution)

Complete the following reaction equations: (3)

* The correct master values will be given to you later,
 values with two digits after the decimal point otherwise -1 point

**2.4 Mass (in mg) of copper in your parent solution,
mass (in mg) of lanthanum in your parent solution. (3)**

$$[M(\text{Cu}) = 63.55 \text{ g mol}^{-1}; M(\text{La}) = 138.91 \text{ g mol}^{-1}]$$

Amount of copper:

$$10,50 \text{ mL} \cdot 0,01 \text{ mol L}^{-1} \cdot 4 \cdot 10 \cdot 63,55 \text{ g mol}^{-1} = 266,9 \text{ mg} \quad (1)$$

Amount of lanthanum:

$$[11,60 - (10,50/10 \cdot 4)] \text{ mL} \cdot 0,1 \text{ mol L}^{-1} \cdot 10 \cdot 138,91 \text{ g mol}^{-1} = 1028 \text{ mg} \quad (2)$$

$$\text{mass Cu} \quad m(\text{Cu}) = 266,9 \text{ mg}$$

$$\text{mass La} \quad m(\text{La}) = 1028 \text{ mg}$$

2.5 Assume a fictive consumption of 39.90 mL of 0.1000 mol L⁻¹ EDTA solution and 35.00 mL of 0.01000 mol L⁻¹ Na₂S₂O₃ solution. Calculate the coefficient x in the formula La_xM_(2-x)CuO₄ (M = Ca and/or Sr and/or Ba) and give the exact formula of the superconductor (5)

$$\text{consumption for lanthanum} = [39,90 - (35,00/10 \cdot 4)] \text{ mL} = 25,90 \text{ mL} \quad (2)$$

$$\text{consumption for copper} = (39,90 - 25,90) \text{ mL} = 14,00 \text{ mL} \quad (2)$$

$$n(\text{La}) : n(\text{Cu}) = 25,90 : 14,00 = 1,85 : 1$$

$$\text{coefficient x: } 1,85 \quad \text{formula: } \text{La}_{1,85}\text{Ba}_{0,15}\text{CuO}_4 \quad (1)$$

2.2 Complexometric Titration

(35.0 points)

$$P = 35 \cdot \left[1 - \frac{|(C1 - (MV1 \cdot PS/100)) - ((MV1 \cdot PS/100) \cdot 0.005)|}{((MV1 \cdot PS/100) \cdot 0.03) - ((MV1 \cdot PS/100) \cdot 0.005)} \right]$$

P = points


C1 = experimental consumption (mL)

MV1 = actual master value

PS = mL of superconductor solution handed out (100.0, 99.00, 98.00, 97.00 mL)

if P ≥ 35 use the maximum points of 35

if P ≤ 0 use zero points



2.3 Iodometric Titration

(35.0 points)

$$P = 35 \cdot \left[1 - \frac{|(C2 - (MV2 \cdot PS/100)) - ((MV2 \cdot PS/100) \cdot 0.0075)|}{((MV2 \cdot PS/100) \cdot 0.04) - ((MV2 \cdot PS/100) \cdot 0.0075)} \right]$$

P = points


C2 = experimental consumption (mL)

MV2 = actual master value

PS = mL of superconductor solution handed out (100.0, 99.00, 98.00, 97.00 mL)

if P ≥ 35 use the maximum points of 35

if P ≤ 0 use zero points



Accuracy of points for 2.2 and 2.3: rounding value is one digit after the decimal point.