


Problem 7



i) Estimate the equilibrium constant for the formation of I_3^- ion from the two curves.


Problem 0 (reserve)

0a	0b	0c	0d	0e	Task 0
4	3	8	8	2	25

The difference in the chemical behaviour of various isotopes is usually negligible, unless the relative change in the molecular mass is considerable.

- a) What would be the highest percentage change in the molecular mass of a neutral molecule upon substitution of a single atom with its isotope?

The following reaction continuously takes place in liquid bromine:



- b) What are the mole fractions of these species in bromine at natural abundance (50 % ${}^{79}\text{Br}$, 50 % ${}^{81}\text{Br}$)?

$x({}^{79}\text{Br}_2)$

$x({}^{81}\text{Br}_2)$

$x({}^{79}\text{Br}-{}^{81}\text{Br})$

- c) Give the equilibrium constant of the process in mole fractions.

$K:$

- d) What is the standard molar entropy change associated with this reaction, supposing that the chemical behaviour of the molecules involved is identical?

$\Delta_r S^\circ:$

Name:

Code: XXX-

In 1913 Hevesy and Paneth carried out the following experiment that began the use of isotopes as tracers and eventually lead to a Nobel Prize for Hevesy for his isotope tracer studies.

They collected in a closed tube a strongly radioactive, completely unreactive gas emitted from radium (at that time known as Curie emanation). It has a relative atomic mass of 222.

e) What is this gas known as today? Give its formula.

This gas was known to go through a succession of radioactive decay processes, forming products called radium A, B, C, D, E, F, G one after the other. The following was already known about this decay chain:

If the gas was left to equilibrate for a few days over water, a solution containing mainly radium D was obtained. After a few weeks, the intensity of the α radiation from Radium F was seen to increase, and slowly reached a steady value, but the quantity of Radium D did not decrease significantly during this time.

f) Which could be the slowest step in the successive transformations?

Radium A \rightarrow B B \rightarrow C C \rightarrow D D \rightarrow E E \rightarrow F F \rightarrow G

Hevesy had earlier worked with Radium D and he was unable to separate it from the inactive lead it was mixed with. The inactive Radium G (atomic mass 206) had shown exactly the same chemical behaviour.

The formation of Radium A, B and G from its precursor was found to be an α -decay, while the formation of Radium C, E and F was a β -decay.

g) How could Radium D form from Radium C? Use α and β in your answer.

In his experiment, Hevesy mixed some aqueous lead-chloride solution (containing 9.69 mg PbCl_2) to the radium D solution he obtained. They measured the β -activity of 1.00 cm^3 of the 120.0 cm^3 mixture to be 16.90 in arbitrary units.

At the same time, with the same equipment they measured the activity of a sample prepared in the following way: Potassium chromate was added to of the lead solution to quantitatively precipitate lead chromate. The precipitate was filtered and was left to stand with approximately 100 cm^3 distilled water for a day at 25 C. The water from this experiment was carefully filtered off. The first part of the filtrate was discarded and 70.0 cm^3 was separated and concentrated by evaporation. Its activity was found to be 0.15 relative units.

Name:

Code: XXX-

h) Give an estimate of the solubility product of lead chromate based on these results.

K_{sp}